



Media Inquiries: Kendra Snyder; Department of Communications
212-496-3419, ksnyder@amnh.org
www.amnh.org

June 27, 2018

STUDY SIGNALS DRAMATIC CHANGE IN WAY ANCIENT DIETS ARE CALCULATED

NEW RESEARCH ON PLANT-EATING MAMMALS FINDS KEY VALUE IN CARBON ISOTOPE ANALYSES IS NOT CONSTANT, CHANGES WITH BODY SIZE

Knowing what extinct animals ate has long been determined by analyzing carbon isotopes locked inside fossil teeth. For two decades, a key isotope value in these equations has been assumed to be the same for all plant-eating mammals, but new research led by Julia Tejada-Lara from the American Museum of Natural History and Columbia University, and published in the journal *Proceedings of the Royal Society B*, contradicts that assumption, revealing that the value changes with body size.

Inferring the diet of extinct vertebrates traditionally depends on anatomical data for a species – for example, the shape of teeth, the height and amount of wear of tooth enamel, and the configuration of the skull and jaw. But more recently, researchers started using stable carbon isotope analyses ($\delta^{13}\text{C}$) to directly determine the foods extinct herbivores ate while alive. Stable carbon isotopes, which form in specific proportions inside plants, are preserved in the body tissues of the animals that eat those plants. By taking samples from the animal's bones, teeth, toenails, or other biological material, paleontologists can determine the kinds of plants that individual animals consumed, how different species might have split up resources in the same habitat, and even overall rainfall or other environmental characteristics.

“Stable carbon isotope studies now are an essential part of our scientific toolkits, informing a wide array of paleontological and geological topics, from predator-prey

interactions to ancient rainfall estimates,” said John Flynn, a co-author on the study and the Frick Curator of Fossil Mammals in the Museum’s Division of Paleontology. “Applying our new formulas that account for body size effects is essential for a more accurate understanding of the history of diverse species, ecosystems, and climates in Earth history.”

As an animal incorporates carbon from its diet into its body tissue, an array of physiological processes affects how much of and in what proportions the carbon isotopes actually get absorbed. Based on studies of cows and other large ungulates, scientists have widely assumed that the skeletal tissues of all plant-eating mammals are “enriched” by about 14 per mil (14 parts in a thousand) in stable carbon isotopes relative to the food that they eat. That constant, called the enrichment value, is an essential part of the equation scientists use for geochemical analyses of animal diets. But Tejada-Lara, a graduate student at the Museum and at the Department of Earth and Environmental Sciences at Columbia University, who was studying sloths, was suspicious that something was overlooked with this one-size-fits-all equation.

“Using a single value for all mammal herbivores, from tiny monkeys to giant elephants, seemed like a large assumption to me,” said Tejada-Lara, the lead author of the study. “I decided to take a closer look at the enrichment value in living and fossil sloths because they are outliers in a number of traits when compared to plant-eating mammals as a whole.”

In collaboration with the Huachipa Zoo in Lima, Peru, Tejada-Lara sampled the food and tooth dust—collected during routine veterinary dental care—of two species of modern sloths: the three-toed sloth (*Bradypus variegatus*) and the two-toed sloth (*Choloepus hoffmanni*). The sloths analyzed were fed a controlled and steady diet from a young age, giving researchers the ability to directly measure the amount of stable carbon isotope they ingest. When the researchers calculated the carbon isotope enrichment value based on samples taken from the sloths’ teeth, they determined it to be 10 per mil for three-toed sloths and 12 per mil for two-toed sloths.

They then measured the enrichment value for the giant extinct ground sloth *Mylodon darwini*, which lived in South America until about 10,000 years ago. By sampling the dung recovered with the fossil and the teeth of the specimen, the researchers determined the tooth tissue was enriched by about 15.5 per mil relative to the proportions in the dietary carbon.

“That’s about a 5-per mil variation just within a single group of mammals,” Tejada-Lara said. “That can make a big difference in the interpretation of research results, including the potential of this discovery to provide more precision on how the evolution of the diet in our own lineage has evolved, because primates include small and large species.”

The researchers combined the sloth data with enrichment values calculated for a range of other mammalian herbivores with known diets, from prairie voles to pigs to giraffes, and found a dramatically clear pattern: the enrichment value generally increases along with body size. They found that at 9.1 per mil, the house mouse (*Mus musculus*) had the lowest value of those measured, and the extinct sloth *Mylodon* had the largest value. Based on this study, the research team has proposed new models of how carbon is incorporated into mammalian plant-eaters, from food to teeth, for the scientific field to integrate into their studies of modern and ancient animals and ecosystems.

The study also found a different pattern of enrichment depending on the makeup of an animal’s digestive tract—specifically whether the animal is a foregut or hindgut fermenter. As it turns out, the giant ground sloth *Mylodon* was a foregut fermenter producing huge amounts of methane.

Other authors on the study include Bruce MacFadden, from the University of Florida; Lizette Bermudez and Gianmarco Rojas from the Huachipa Zoo; and Rodolfo Salas-Gismondi from the Cayetano Heredia University and the San Marcos National University in Peru.

Funding was provided by the Florida Museum of Natural History, U.S. National Science Foundation grant# PIRE 0966884, Columbia University’s Graduate School of Arts

and Sciences, and the American Museum of Natural History's Division of Paleontology Frick Fund.

Proceedings of the Royal Society B paper:

<http://rspb.royalsocietypublishing.org/content/285/1881/20181020>

AMERICAN MUSEUM OF NATURAL HISTORY (AMNH.ORG)

The American Museum of Natural History, founded in 1869, is one of the world's preeminent scientific, educational, and cultural institutions. The Museum encompasses 45 permanent exhibition halls, including those in the Rose Center for Earth and Space and the Hayden Planetarium, as well as galleries for temporary exhibitions. It is home to the Theodore Roosevelt Memorial, New York State's official memorial to its 33rd governor and the nation's 26th president, and a tribute to Roosevelt's enduring legacy of conservation. The Museum's five active research divisions and three cross-disciplinary centers support approximately 200 scientists, whose work draws on a world-class permanent collection of more than 34 million specimens and artifacts, as well as specialized collections for frozen tissue and genomic and astrophysical data, and one of the largest natural history libraries in the world. Through its Richard Gilder Graduate School, it is the only American museum authorized to grant the Ph.D. degree. Beginning in 2015, the Richard Gilder Graduate School also began granting the Master of Arts in Teaching (MAT) degree, the only such freestanding museum program. Annual visitation has grown to approximately 5 million, and the Museum's exhibitions and Space Shows are seen by millions more in venues on six continents. The Museum's website, mobile apps, and MOOCs (massive open online courses) extend its scientific research and collections, exhibitions, and educational programs to additional audiences around the globe. Visit amnh.org for more information.

Follow

Become a fan of the Museum on Facebook at [facebook.com/naturalhistory](https://www.facebook.com/naturalhistory), follow us on Twitter at @AMNH, on Instagram at [@AMNH](https://www.instagram.com/AMNH), and on Tumblr at amnhnyc.

#

No. 44

(more)

4